


The Orthopaedic and Fracture Clinic 1431 Premier Drive Mankato, MN 56001 507-386-6600

Dr. Allen Ganglion Cyst Removal Post-Op Protocol

Post-Op Instructions

- 1. Keep dressings in place until post op day #3
- 2. Keep hand and dressings clean and dry, cover them to shower and bathe
- 3. Ok to get wound wet in shower post-op day #5, no soaking until after first post-op visit (2 weeks)
- 4. Cryotherapy with ice or a cool pack several times a day
- 5. Scheduled Tylenol and Ibuprofen for pain control
 - a. Limited narcotics are given for cyst removal, this is to be used for break through pain only
- Local anesthetic is injected at the time of the surgery for pain control, this will last 4-8 hours
- 7. Keep hand elevated to improve swelling into your fingers
- 8. Work on elbow, wrist, hand, and finger ROM



STRAIGHT HOOK FULL FIST TABLE TOP ST

- 9. No heavy lifting (<3# limit)
- 10. Begin tendon & nerve glide exercises immediately post-op

First Post-Op Visit (~POD #14)

- 1. Follow up from surgery at 2 weeks post-op
- 2. Sutures will be removed at this appointment
- 3. Continue to work on hand and finger motion with emphasis on tendon and median nerve glides
- 4. Hand strengthening with putty or a Theraball at 3-4 weeks post-op

Second Post-Op Visit (POD~28 [OPTIONAL])

1. Depending on your progress you may elect to cancel this appointment

Recovery

- 1. You may do light work such as typing within the first week after surgery
- 2. Return to strenuous work can take 4-6 weeks
- 3. Pinch strength returns in 2-6 weeks
- 4. Grip strength returns in 6-12 weeks