

## Dr. Allen Triceps Repair Post-Op Instructions

### Dressings

Keep the splint in place and your surgical dressings in place until the first post-op visit at 2 weeks. Use a bag over your arm or take sponge baths during this time. After your first visit your splint will be removed and exchanged for a brace. This brace can be removed for hygiene. Make sure to wash your hands before touching the skin around your incision. Dressings can be removed and replaced with clean, fresh gauze, make sure to keep the wound covered until your first follow up. It is ok to shower at 2-weeks post-op. Allow clean water to wash over the wound, do not scrub or wash it with any soaps. Your wound was likely closed with absorbable sutures and glue. Do not pick at or remove any scabs or loose glue ends, they will fall off on their own. It is ok to lotion the skin surrounding the incision, but don't place any lotion or ointment directly on the wound. Once you've showered and replaced the dressings, replace your brace. Avoid any flexing of your elbow past 90° during this time.

### Brace

Your motion will be restricted for the first 6 weeks after surgery to protect the repair. Your brace will be adjusted at your weekly PT visits. The exact timing of regaining motion is individualized based on the tension of the repair at the time of surgery. Generally, it will be no motion for 2 weeks, then gradually increasing motion over the next 4 weeks until full motion is regained around 6 weeks. It is important to not exceed the motion limited to protect the repair while it heals.

### Pain Control

Pain control is a very important aspect of your recovery. It is well documented that multimodal pain control is the most effective way to manage post-operative pain. At the end of the procedure, I will inject local anesthetic into the surgical field, additionally the anesthesiologist may give you an upper extremity block. These blocks can last anywhere for a few hours to a couple days. The goal is to have little to no pain in the immediate post-operative period. You will also receive a prescription for narcotic pain medications. This is to be used for break through pain. Combining the narcotic pain medications with Tylenol and Ibuprofen has been shown to give superior pain relief

compared to the narcotics alone. The goal is to use the narcotic pain medications sparingly, and discontinue its use by 1-week post-op. If you are running low and will require a refill, please plan in advance, as refills on the weekend are often not logistically possible. Ice and elevation for the first several days after your surgery will also help control your pain.

### **Protection**

The triceps is an extremely strong muscle, and you can easily damage or injure the repair during the rehabilitation process. It is essential to not push yourself up from a chair, do any heavy lifting, or have any sudden jerking movements with the operative arm until cleared by Dr. Allen and PT. This is to protect the repair while it heals.

### **Nausea**

Nausea after surgery and anesthesia is not uncommon. You will receive an anti-nausea medication prescription for use at your discretion.

### **Constipation**

Narcotic pain medications can be very constipating. You will receive a stool softening medication. I recommend you take it as long as you are taking the narcotics. Make sure to drink plenty of water as well in the immediate post-operative period.

### **Physical Therapy**

Your first PT visit will 2 weeks after your surgery. Therapy is a critical aspect of regaining function, motion, and strength after triceps repair.

### **Follow Up**

Typically, your first follow up is about 2 weeks after surgery. We will assess your wound and remove any sutures at this time. We will remove your splints, place the elbow brace, and set you up with your first PT visit after this appointment as well.